

AD-A148 382 MODULATIONAL INSTABILITY OF LOWER HYBRID WAVES(U) REGIS 1/1
COLL RESEARCH CENTER WESTON MA S P KUO ET AL.
01 JUN 84 AFGL-TR-84-0302 F19628-83-K-0024

UCLASSIFIED


F/G 12/1

NL

END

FILMED

DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS None	
2a. SECURITY CLASSIFICATION AUTHORITY N/A		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release, distribution unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE N/A		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFGL-TR-84-0302	
4. PERFORMING ORGANIZATION REPORT NUMBER(S)		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFGL-TR-84-0302	
6a. NAME OF PERFORMING ORGANIZATION Regis College	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Air Force Geophysics Laboratory/LIS	
6c. ADDRESS (City, State and ZIP Code) 235 Wellesley Street Weston, Mass. 02193		7b. ADDRESS (City, State and ZIP Code) Hanscom AFB, MA 01731	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Air Force Geophysics Lab	8b. OFFICE SYMBOL (If applicable) LIS	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F19628-83-K-0024	
8c. ADDRESS (City, State and ZIP Code) Hanscom AFB, MA 01731 John A. Klobuchar		10. SOURCE OF FUNDING NOS.	
		PROGRAM ELEMENT NO. 62101F	PROJECT NO. 4643
		TASK NO. 09	WORK UNIT NO. AB
11. TITLE (Include Security Classification) Modulational instability of lower hybrid waves			
12. PERSONAL AUTHOR(S) S.P. Kuo, M.C. Lee			
13a. TYPE OF REPORT Scientific - Reprint	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Yr., Mo., Day) June 1, 1984	15. PAGE COUNT 4
16. SUPPLEMENTARY NOTATION *Polytechnic Institute of NY, Long Island Center, Farmingdale, NY 11735 Proceedings of 1984 International Conference on Plasma Physics, Lausanne, Switzerland, June 27 - July 3, 1984			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB GR.	
		Modulational instability, lower hybrid waves; Purely growing modes	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) We investigate the excitation of a purely growing modulational mode together with two lower hybrid sidebands by a lower hybrid pump. The condition has been derived. The growth rate of the instability is found to be quite large for the power density employed in the lower hybrid heating experiments in large tokamaks.			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL John A. Klobuchar		22b. TELEPHONE NUMBER (Include Area Code) 617-861-3988	22c. OFFICE SYMBOL AFGL/LIS

DD FORM 1473, 83 APR

EDITION OF 1 JAN 73 IS OBSOLETE.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

AD-A148 382

DTIC
ELECTE
DEC 7 1984

AFGL-TR-84-0302

Modulational instability of lower hybrid waves

S.P. Kuo

Polytechnic Institute of New York, Long Island Center,
Farmingdale, N.Y. 11735, U.S.A.

M.C. Lee

Regis College Research Center, Weston, Mass. 02193, U.S.A.

Abstract. We investigate the excitation of a purely growing modulational mode together with two lower hybrid sidebands by a lower hybrid pump. The condition, $|k_{oz}/k_{ol}| < \omega_{pi}/\Omega_e$, needs to be satisfied for the instability, where $|k_{oz}/k_{ol}|$ is the ratio of the perpendicular to the parallel scale lengths of the lower hybrid pump. The growth rate of the instability is found to be quite large for the power densities employed in the lower hybrid heating experiments in large tokamaks.

1. Introduction

The need of supplementary plasma heating in magnetic fusion devices has been generally recognized for achieving the operational temperatures for fusion power production. Among various RF heating schemes, lower hybrid waves have received much attention not only for plasma heating (e.g., Porklab, 1977) but also for plasma confinement (e.g., Fisch, 1978) purposes. The effectiveness of these processes relies on the penetration of RF wave energy into the interior of the plasmas. Hence, the self-modulational effects on the lower hybrid wave propagation are believed to be important on this aspect (e.g., Morales and Lee, 1975). In this paper, we perform a linear stability analyses of the modulational instability whereby a purely growing modulational mode together with two lower hybrid sidebands are excited by a lower hybrid pump wave.

2. Coupled mode equations

We consider the propagation of a lower hybrid pump wave $\phi = \tilde{\phi}_0 \exp[i(\mathbf{k}_0 \cdot \mathbf{r} - \omega_0 t)]$ in a uniform, collisionless plasma embedded in a constant magnetic field $\mathbf{B}_0 = \hat{z}B_0$. Here, ϕ_0 is the lower hybrid wave field potential; $\mathbf{k}_0 (= \hat{y}k_{ol} + \hat{z}k_{oz})$ and ω_0 are respectively, the wave vector and the angular wave frequency of the lower hybrid pump that satisfies the dispersion relation: $\omega_0 = \omega_{LH}[1 + (M/m)(k_{oz}/k_{ol})^2]^{1/2}$, where $\omega_{LH} = \omega_{pi}/(1 + \omega_{pe}^2/\Omega_e^2)^{1/2}$ is the lower hybrid resonance frequency.

Proceedings of 1984 International Conference on Plasma Physics, Lausanne, Switzerland, June 27 - July 3, 1984

84 11 27 040

The parameters, M/m , ω_{pi}/ω_{pe} , and Ω_e are the ratio of ion to electron masses, the ion(electron) plasma frequency, and the electron gyro-frequency, respectively.

The process under consideration is a modulational instability whereby the lower hybrid pump wave (ω_0, k_0) excites two lower hybrid sidebands (ω_{\pm}, k_{\pm}) and a purely growing modulational mode (ω_s, k). This process can be described by the following frequency and wave vector matching relations: $\omega_+ - \omega_s = \omega_0 = \omega_- + \omega_s^*$ and $k_{\pm} = k_0 \pm k$, where $k \parallel k_0$. The lower hybrid sidebands are excited through the beating current density driven by the lower hybrid pump field on the density perturbations, $n_s = \tilde{n}_s \exp[i(k_s \cdot r - \omega_s t)]$, of the zero-frequency modulational mode.

From the Poisson's equation, $k_{\pm}^2 \phi_{\pm} = 4\pi e(\delta n_{i\pm} - \delta n_{e\pm})$, where the ion and electron density perturbations ($\delta n_{i\pm}$ and $\delta n_{e\pm}$) associated with the lower hybrid sidebands are given, respectively, by $\delta n_{i\pm} = (n_0 e / M \omega_0^2) k_{\pm}^2 [\phi_{\pm} + (k_{01}/k_{\pm 1})(\omega_0/\omega_{\pm})\phi_0(n_{s\pm}/n_0)]$ and $\delta n_{e\pm} = (n_0 e / m \Omega_e^2)(1 - k_{0z}^2 \Omega_e^2 / k_{01}^2 \omega_0^2) k_{\pm 1} [\phi_{\pm} + (k_{01}/k_{\pm 1})(\omega_0/\omega_{\pm})\phi_0(n_{s\pm}/n_0)]$, the coupled mode equation for the lower hybrid sidebands is obtained as

$$\phi_{\pm} = (k_0/k_{\pm})(\omega_{\pm}/\omega_0)(k_{\pm 1}^2/k_{\pm}^2)[\omega_{LH}^2 \omega_0^2 / \omega_{pi}^2 \omega_{s\pm}(\omega_{s\pm} \pm 2\omega_0)]\phi_0(n_{s\pm}/n_0) \quad (1)$$

where the relations $\omega_{s+} = \omega_s = \omega_s^*$ have been used.

The nonlinear effects responsible for the excitation of the purely growing modulational mode include the nonlinear beating current and the pondermotive force appearing, respectively, in the continuity equation and the momentum equation. Under the assumption of quasi-neutrality, the coupled mode equation for the purely growing mode is derived from the continuity and the momentum equations of electrons and ions and has the following form:

$$\begin{aligned} [-\omega_s^2(1 + k_{is}^2 c_s^2 / \Omega_e \Omega_i) + k_{zs}^2 c_s^2](n_s/n_0) = & -\omega_s k_{1F}(\delta v_{ey}^{NL} + \frac{m}{M} \delta v_{iy}^{NL}) \\ & - k_{zs}(\delta v_{ez}^{NL} + \frac{m}{M} \delta v_{iz}^{NL}) + k_{1s} \frac{\omega_s}{\Omega_e} (\frac{F_{ix}}{M} - \frac{F_{ex}}{m}) + ik_{1s} \frac{\omega_s^2}{m \Omega_e^2} (\\ & F_{iy} + F_{ey}) - i \frac{k_z}{M} (F_{iz} + F_{ez}) \end{aligned} \quad (2)$$

where $C_s = [(T_e + T_i)/M]^{1/2}$ is the ion acoustic velocity. The nonlinear beating current density $n_0 \delta v_{e,i}^{NL}$ and the pondermotive force $F_{e,i}$ are given, respectively, by

$$n_o \delta v_{e,i}^{NL} = \delta n_{e,i0} \delta v_{e,i-}^* + \delta n_{e,i0}^* \delta v_{e,i+} + \delta n_{e,i-}^* \delta v_{e,i0} + \delta n_{e,i+} \delta v_{e,i0}^* \\ \cdot \delta v_{e,i0}^*$$

$$\text{and } F_{e,i} = -im_{e,i} \{ k_o \cdot \delta v_{e,i-}^* \delta v_{e,i0} - k_- \cdot \delta v_{e,i0} \delta v_{e,i-}^* + k_+ \delta v_{e,i0}^* \delta v_{e,i+} \\ \cdot \delta v_{e,i+} - k_o \cdot \delta v_{e,i+} \delta v_{e,i0}^* \}$$

$$\text{where } \delta v_{e,\alpha} = -ik_{\alpha 1} [\hat{x} + i\hat{y}(\omega_{\alpha}/\Omega_e) - i\hat{z}(k_{ox}\Omega_e/k_{o1}\omega_{\alpha})] (e\phi_{\alpha}/m\Omega_e),$$

$$\alpha = 0 \text{ or } \pm \quad (3)$$

$$\delta v_{i0} = ik_{o1} [\hat{x}(\Omega_i/\Omega_o) - i\hat{y} - i\hat{z}(k_{oz}/k_{o1})] (e\phi_o/M\omega_o), \quad (4)$$

$$\delta v_{i\pm} = ik_{\pm 1} [\hat{x}(\Omega_i^2/\omega_{\pm}^2) - i\hat{y}(\Omega_i/\omega_{\pm}) - i\hat{z}(k_{oz}\Omega_i/k_{o1}\omega_{\pm})] (e\phi_{\pm}/m\Omega_e), \quad (5)$$

$$\delta n_{eo} = n_o k_{o1}^2 (1 - k_{oz}^2 \Omega_e^2 / k_{o1}^2 \omega_o^2) (e\phi_o / m\Omega_e^2) \quad (6)$$

$$\delta n_{io} = n_o k_o^2 (e\phi_o / M\omega_o^2). \quad (7)$$

With the aid of (3) - (7), two things have been noted from the RHS of (2): Firstly, the ion-nonlinearity terms are relatively insignificant compared to the electron-nonlinearity terms; Secondly, the nonlinear beating current effect is partially cancelled by the pondermotive force effect. Therefore, the electron-nonlinearity term that remains on the RHS of (2) is $-\omega_s k_z \delta v_{ez}^{NL}$, contributed from the induced nonlinear beating current flowing along the imposed magnetic field.

Eliminating n_s and ϕ_{\pm} from Equations (1) and (2) leads to the dispersion relation

$$\omega_s^2 = k_z^2 \{ C_s^2 - [(1 - k_{oz}^2 \Omega_e^2 / \omega_{pi}^2 k_{o1}^2) \Omega_e^2 / (1 + Mk_{oz}^2 / mk_{o1}^2) (\Omega_e^2 + \omega_{pe}^2)] \\ \cdot |v_{\phi}|^2 \} / (1 + k_1^2 C_s^2 / \Omega_e \Omega_i) \quad (8)$$

where $v_{\phi} = k_o e\phi_o / m\Omega_e$. For a purely growing mode (i.e., $\omega_s = i\gamma$), the growth rate obtained from (8) is given by

$$\gamma = k_z \{ [(1 - k_{oz}^2 \Omega_e^2 / \omega_{pi}^2 k_{o1}^2) \Omega_e^2 / (1 + Mk_{oz}^2 / mk_{oz}^2) (\Omega_e^2 + \omega_{pe}^2)] |v_{\phi}|^2 - C_s \}^{1/2} / (1 + k_1^2 C_s^2 / \Omega_e \Omega_i)^{1/2}. \quad (9)$$

Since γ has to be positive for instability to occur, it is clear from (9) that (I) $|k_{oz}/k_{o1}| < \omega_{pi} \Omega_e$, (10)
namely, the ratio of the perpendicular to the parallel scale lengths of the lower hybrid pump wave has to be less than the ratio of the ion plasma frequency to the electron gyrofrequency, (II) $|\tilde{\phi}_o| > |\tilde{\phi}_{oth}| = ($

$$m/e)(\Omega_e^2 + \omega_{pe}^2)^{1/2}(C_s/k_o)(1 + Mk_{oz}^2/mk_{ol}^2)^{1/2}/(1 - \Omega_e^2 k_{oz}^2/\omega_{pi}^2 k_{ol}^2)^{1/2} \quad (11)$$

namely, the pump field intensity represented by $|\tilde{\phi}_0|$ has to be greater than a threshold defined by $|\tilde{\phi}_{oth}|$.

The growth rate has a minimum when $k_{oz}^2 \Omega_e^2/\omega_{pi}^2 k_{ol}^2 = (1 - |\tilde{\phi}_{oth}/\tilde{\phi}_0|^2)/(2 - |\tilde{\phi}_{oth}/\tilde{\phi}_0|^2)$. In this optimum case and for a strong lower hybrid pump wave, (9) may be expressed as

$$\gamma_m \sim k_z |v_\phi| \Omega_e^2/(2\Omega_e^2 + \omega_{pe}^2)^{1/2}(\Omega_e^2 + \omega_{pe}^2)^{1/2}(1 + k_i^2 C_s^2/\Omega_e \Omega_i)^{1/2} \quad (12)$$

which is proportional to the wave length of the modulational instability.

4. Summary and conclusions

The condition and the threshold power have been derived for the modulational instability of lower hybrid waves as shown, respectively, in (10) and (11). The nonlinearity for the excitation of purely growing modes is provided by the nonoscillatory beating current in the direction of the imposed magnetic field. For the power densities commonly used in the lower hybrid heating experiments in large tokamaks, the growth rate given in (12) is found to be quite large and it covers a broad spectrum. However, it should be stressed that the spatial modulational on the lower hybrid pump wave as discussed here only occurs in the resonance region where the condition, $|k_{oz}/k_{ol}| < \omega_{pi}/\Omega_e$, can be satisfied.

References

- Fisch, N.J., Phys. Rev. Lett. 41, 873, 1978.
 Morales, G.J., and Y.C. Lee, Phy. Rev. Lett., 35, 930, 1975.
 Porklab, M., Phys. Fluids, 20, 2058, 1977.

Acknowledgments. This work was supported jointly in part by the NSF grant ATM-8315322 and in part by the AFSOR grant AFSOF-83-0001 at Polytechnic Institute of New York and by AFGL contract F19628-83-K-0024 at Regis College Research Center.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A/1	1



END

FILMED

1-85

DTIC